

Circular Economy in Food Industry Waste Management (BIVALIA Workshop2 - 2026)

Editors: Elena Díaz Nieto; Francisco Heras Muñoz; Juan Antonio Melero Hernández; Manuel Gertrudix
Madrid, BIVALIA, UAM-URJC, 2026

DOI: 10.5281/zenodo.22206833 · CC BY 4.0

Full volume: <https://books.ciberimaginario.cloud/libro/bivalia-w2-2026>

OFFPRINT · P. 40

P.I.5 — Growth and tolerance screening of engineered *Shimwellia blattae* and *Escherichia coli* on agro-industrial residues for isobutanol production

C. López-Martínez¹, M.J. Negro¹, I. Ballesteros¹, A.D. Moreno¹, M.G. Acedos^{1*}

1: Advanced Biofuels and Bioproducts Unit, Energy Department, CIEMAT, Avda. Complutense 40, 28040 Madrid, Spain

Isobutanol production; Agro-industrial residues; Engineered bacteria; Growth and tolerance screening

HOW TO CITE THIS CONTRIBUTION

C. López-Martínez, M.J. Negro, I. Ballesteros, A.D. Moreno, M.G. Acedos. (2026). P.I.5 — Growth and tolerance screening of engineered *Shimwellia blattae* and *Escherichia coli* on agro-industrial residues for isobutanol production. In: Díaz Nieto, E., Heras Muñoz, F., Melero Hernández, J. A., Gertrudix, M. (Eds.), Circular Economy in Food Industry Waste Management (BIVALIA Workshop2 - 2026), p. 40. BIVALIA, UAM-URJC. <https://doi.org/10.5281/zenodo.22206833>

Poster Presentations #22

P.I.5 — Growth and tolerance screening of engineered *Shimwellia blattae* and *Escherichia coli* on agro-industrial residues for isobutanol production

C. López-Martínez¹, M.J. Negro¹, I. Ballesteros¹, A.D. Moreno¹, M.G. Acedos^{1*}

¹ Advanced Biofuels and Bioproducts Unit, Energy Department, CIEMAT, Avda. Complutense 40, 28040 Madrid, Spain

*Correspondence: miguel.garciaacedos@ciemat.es

Keywords: Isobutanol production; Agro-industrial residues; Engineered bacteria; Growth and tolerance screening

Biorefineries are driving interest in transforming agro-industrial residues into valuable products and biofuels. Among these, isobutanol has emerged as a particularly attractive alternative to ethanol due to its higher energy density, lower hygroscopicity, and improved compatibility with existing fuel infrastructure. The sustainable production of biofuels from agro-industrial residues therefore represents a promising strategy to reduce reliance on fossil resources while valorizing waste streams. In this study, the potential of two genetically engineered bacterial strains, *Shimwellia blattae* p424IbPSO and *Escherichia coli* W pIZIbPSO [1, 2], for isobutanol production using agro-industrial residues was evaluated. A preliminary screening was performed to assess microbial growth and tolerance to different waste-derived substrates. The tested feedstocks included cheese whey from the dairy industry, brewer's spent grain (BSG) from the brewing process, and a hot water extract of sugars obtained from grape stems generated in the wine industry. These residues were incorporated at different concentrations into a defined minimal salts medium (M92X), which served as the base culture medium. Growth performance and substrate compatibility were compared against control cultures grown in standard M92X medium. Cultures were conducted in multiwell plates under controlled conditions in a plate reader CLARIOstar (BMG Labtech), with the addition of selective antibiotics (gentamicin and streptomycin) to ensure plasmid maintenance. The screening results demonstrated that both strains were capable of growing on tested residues, although growth profiles and tolerance thresholds varied depending on the substrate type and concentration. Cheese whey and grape stem extracts supported robust growth at moderate concentrations, while higher loadings of brewer's spent grain-derived substrates showed inhibitory effects, likely due to complex matrix components. These findings highlight the adaptability of the engineered strains to heterogeneous, low-cost substrates and provide key insights into optimal substrate concentrations for subsequent fermentation processes. The results will guide future optimization of isobutanol production using agro-industrial waste streams as feedstock.

Acknowledgements

This work was carried out within the framework of the BIVALIA-CM project [TEC-2024/BIO-177], funded by the Technology Networks call of the Community of Madrid. The authors thank Lácteas del Jarama, Cervezas La Cibeles, and Bodega Bernabeleva for providing the raw materials used in this study. The authors also acknowledge José Luis García López and Beatriz Galán (Centro de Investigaciones Biológicas Margarita Salas, CSIC) for kindly providing the microbial strains.

References

- [1] C. Felpeto-Santero, A. Rojas, M. Tortajada, B. Galán, D. Ramón, and J. L. García, "Engineering alternative isobutanol production platforms," *AMB Express*, vol. 5, no. 1, p. 32, 2015. <https://doi.org/10.1186/s13568-015-0119-2>
- [2] M. G. Acedos, I. De la Torre, V. E. Santos, F. García-Ochoa, J. L. García, and B. Galán, "Modulating redox metabolism to improve isobutanol production in *Shimwellia blattae*," *Biotechnology for Biofuels*, vol. 14, no. 1, p. 8, 2021. <https://doi.org/10.1186/s13068-020-01862-1>